



King's Research Portal

DOI:

[10.1016/j.jid.2017.05.014](https://doi.org/10.1016/j.jid.2017.05.014)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Spuls, P. I., Gerbens, L. A. A., Apfelbacher, C. J., Wall, D., Arents, B. W. M., Barbarot, S., Roberts, A., Deleuran, M., Middelkamp-Hup, M. A., Vestergaard, C., Weidinger, S., Schmitt, J., Irvine, A. D., & Flohr, C. (2017). The international TREATment of ATopic eczema (TREAT) Registry Taskforce: an initiative to harmonise data collection across national atopic eczema photo- and systemic therapy registries. *Journal of Investigative Dermatology*, 2014-2016. <https://doi.org/10.1016/j.jid.2017.05.014>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

The international TREatment of ATopic eczema (TREAT) Registry Taskforce: an initiative to harmonise data collection across national atopic eczema photo- and systemic therapy registries

Phyllis I. Spuls, MD, PhD, Louise A.A. Gerbens, MD, Christian J. Apfelbacher, MSc, PhD, Dmitri Wall, MD, MSc, Bernd W.M. Arents, Sebastien Barbarot, MD, PhD, Amanda Roberts, Mette Deleuran, MD, PhD, Maritza A. Middelkamp-Hup, MD, PhD, Christian Vestergaard, MD, PhD, Stephan Weidinger, MD, PhD, Jochen Schmitt, MD, Alan D. Irvine, MD, DSc, Carsten Flohr, MD, PhD

PII: S0022-202X(17)31559-2

DOI: [10.1016/j.jid.2017.05.014](https://doi.org/10.1016/j.jid.2017.05.014)

Reference: JID 891

To appear in: *The Journal of Investigative Dermatology*

Received Date: 21 March 2017

Revised Date: 8 May 2017

Accepted Date: 9 May 2017

Please cite this article as: Spuls PI, Gerbens LAA, Apfelbacher CJ, Wall D, Arents BWM, Barbarot S, Roberts A, Deleuran M, Middelkamp-Hup MA, Vestergaard C, Weidinger S, Schmitt J, Irvine AD, Flohr C, The international TREatment of ATopic eczema (TREAT) Registry Taskforce: an initiative to harmonise data collection across national atopic eczema photo- and systemic therapy registries, *The Journal of Investigative Dermatology* (2017), doi: 10.1016/j.jid.2017.05.014.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



The international TReatment of ATopic eczema (TREAT) Registry

Taskforce: an initiative to harmonise data collection across national atopic eczema photo- and systemic therapy registries

Phyllis I. Spuls, MD, PhD,^{1*} Louise A.A. Gerbens, MD,^{1*} Christian J. Apfelbacher, MSc, PhD,² Dmitri Wall, MD, MSc,^{3,4} Bernd W.M. Arents,⁵ Sebastien Barbarot, MD, PhD,⁶ Amanda Roberts,⁷ Mette Deleuran, MD, PhD,⁸ Maritza A. Middelkamp-Hup, MD, PhD,¹ Christian Vestergaard, MD, PhD,⁸ Stephan Weidinger, MD, PhD,⁹ Jochen Schmitt, MD,^{10,11} Alan D. Irvine, MD, DSc,^{3,12,13} and Carsten Flohr, MD, PhD¹⁴

¹ Department of Dermatology, Academic Medical Centre, University of Amsterdam, Amsterdam, The Netherlands

² Medical Sociology, Institute of Epidemiology and Preventive Medicine, University of Regensburg, Regensburg, Germany

³ Department of Paediatric Dermatology, Our Lady's Children's Hospital, Crumlin, Dublin, Ireland

⁴ Irish Skin Foundation, Dublin, Ireland

⁵ Dutch society of patients with atopic eczema (VMCE, Vereniging voor Mensen met Constitutioneel Eczeem), Nijkerk, The Netherlands

⁶ Department of Dermatology, Nantes University Hospital, Nantes, France

⁷ Nottingham Support Group for Carers of Children with Eczema, Nottingham, UK

⁸ Department of Dermatology and Venereology, Aarhus University Hospital, Aarhus, Denmark

⁹ Department of Dermatology and Allergy, University Hospital Schleswig-Holstein, Campus Kiel, Kiel, Germany

¹⁰ Center for Evidence-based Healthcare, Medizinische Fakultät Carl Gustav Carus, TU Dresden, Dresden, Germany

¹¹ University Allergy Center, University Hospital Carl Gustav Carus Dresden, Germany

¹² Department of Clinical Medicine, Trinity College Dublin, Dublin, Ireland

¹³ National Children's Research Centre, Dublin, Ireland

¹⁴ Unit for Population-Based Dermatology Research, St John's Institute of Dermatology, Guy's & St Thomas' NHS Foundation Trust and King's College London, London, UK

* Both authors contributed equally and share first authorship

Corresponding author:

Phyllis I. Spuls

Department of Dermatology, Academic Medical Centre, University of Amsterdam, The Netherlands

Meibergdreef 9

1105 AZ Amsterdam, The Netherlands

Phone: +31 20 5664763

Fax: + 31 20 5669152

E-mail: ph.i.spuls@amc.uva.nl

ABBREVIATIONS

AE: Atopic eczema

EMA: European Medicines Agency

ETFAD: European Taskforce for Atopic Dermatitis

FDA: Food and Drug Administration

IEC: International Eczema Council

ISAD: International Society for Atopic Dermatitis

PARENT JA: PATient REGistries iNiTiative joint action

TREAT: TREatment of ATopic eczema

KEYWORDS

Atopic eczema, Atopic dermatitis, Patient registries, Immuno-modulatory therapies,
Phototherapy

TO THE EDITOR

There is an urgent need for novel immuno-modulatory treatments for patients with atopic eczema (AE) (syn. 'atopic dermatitis'); in particular for those with disease recalcitrant to topical therapies. The only conventional systemic therapy approved by the European Medicines Agency (EMA) is cyclosporine; and only in adults. In the US, only oral corticosteroids and dupilumab have Food and Drug Administration (FDA) approval.

At present, the main evidence to inform clinical practice around conventional systemic immuno-modulatory treatments is derived from a rather small body of randomised controlled trials (RCTs) (Roekevisch et al., 2014) as well as case series (i.a. (Garritsen et al., 2015; van der Schaft et al., 2015)). Despite this, in the absence of approved alternatives, immuno-modulatory treatments are frequently prescribed as off-label therapies in children and adults, as shown in our recent European and North American surveys (Proudfoot et al., 2013; Taylor et al., 2016; Totri et al., 2017). Inclusion criteria for clinical trials are stringent; research in psoriasis suggests that around 30% of patients on systemic therapies entered into registries (in itself a very selected subgroup of all patients on systemic therapy) would not be eligible for a clinical trial, underscoring the value of 'real world' patient data (Garcia-Doval et al., 2012).

With the dawn of a new therapeutic era of biologic therapies for AE, some of which have shown great promise in placebo-controlled studies and have been approved by the FDA (Beck et al., 2014; Simpson et al., 2016; Thaci et al., 2016), we need more evidence with regard to effectiveness, safety and cost-effectiveness of such therapies compared to currently used treatment modalities, such as phototherapy and systemic immuno-modulatory drugs (cyclosporine, methotrexate, azathioprine and others).

Apart from the need for comparative real life clinical data of conventional and new therapies, clinical decision-making requires long-term follow-up data on such treatments to generate

information on disease control/disease trajectory modification (even after treatment has been discontinued) and drug safety, including rare adverse events. Generation of reliable data to address these issues requires observation of large patient cohorts for several years. Comprehensive health economic evaluation of these different treatments is routinely required by many national health technology assessment organizations and third party payers to inform the allocation of health care resources. Consequently, independent prospective multi-centre registries are a logical step, as has been successfully developed in psoriasis. This is all the more important in the light of emerging evidence that suggests that industry-funded post-marketing studies, which are often conducted to detect rare adverse events, are not improving drug safety surveillance (Spelsberg et al., 2017).

The international TREatment of ATopic eczema (TREAT) Registry Taskforce (www.comet-initiative.org/studies/details/825?result=true) seeks to find consensus on core domains and domain items for AE research registries and to harmonise data collection on patients receiving systemic immuno-modulatory therapies, following recently published best-practice guidelines of the European Commission funded PATient REGistries iNiTiative joint action (PARENT JA) (<http://patientregistrieseu/guidelines>) (Zaletel and Kralj, 2015). PARENT was a collaborating partner of the TREAT Registry Taskforce. This approach will reduce heterogeneity between national registries, enhance interoperability of the national registries, allow direct comparability of individual country data and facilitate data pooling between countries. It will also help to establish compatible data entry platforms.

As a first step on this journey, we recently conducted an international eDelphi exercise and a consensus meeting among 410 dermatologists, nurses, non-clinical researchers involved in AE, pharmaceutical industry representatives, regulatory body representatives and patients from 36 countries, including members of the International Eczema Council (IEC), the European Taskforce for Atopic Dermatitis (ETFAD), and the International Society for Atopic

Dermatitis (ISAD). The study identified a core set of domains and domain items to be captured by national AE patient registries (the ‘What’ to measure) (Table 1) (Gerbens et al., 2017). Further work is currently underway to define how these domains and domain items should be measured (the ‘How’ to measure).

We envisage that eligible patients will be children and adults diagnosed with AE who are starting on photo- or systemic immuno-modulatory therapy (conventional immuno-suppressive therapies as well as new biologics), taking into account the national and local eligibility criteria for these specific therapies. Patients receiving intensive topical therapy or screen failures for systemic therapy may also be included. Patients will be followed independently of stopping therapy or subsequent switch to other therapies.

We look forward to working with the global dermatology community on this project, especially those who routinely treat patients with recalcitrant AE; a disease that accounts for more than 20% of total health loss due to skin conditions (Hay et al., 2014) and that has shown a health economic impact similar to that of diabetes mellitus and asthma (Williams et al., 2008). Ultimately, national cohorts of AE patients on photo- and systemic immuno-modulatory therapies will inform treatment guidelines and will also act as a resource for biomarker discovery and pharmacogenetic and pharmacodynamic research. With novel biologic therapies soon to enter our clinical practice, the timing for this project could not be better.

CONFLICT OF INTEREST

AI has served as a consultant to Anacor, Chugai Pharma, Pfizer, Regeneron, Roche/Genentech, Sanofi-Genzyme and UCB Pharma. CV has advised and given lectures for AbbVie, Chugai, Novartis, Regeneron, Sanofi-Genzyme and UCD Pharma. JS has the lead over the German atopic eczema registry (TREAT Germany) and has received institutional funding for investigator-initiated research from ALK, MSD, Novartis, Pfizer, Sanofi and Wyeth. MD has been a speaker, advisory board member and (principal) investigator for AbbVie, CK-Care Foundation, La Roche Posay Laboratoire Dermatologique, Leo Pharma, Meda Pharma, Pierre Fabre Laboratories, Regeneron, Roche and Sanofi-Genzyme. PS has served as a consultant to AbbVie, Anacor, Leo Pharma and Novartis, has received independent research grants from Leo Pharma and Schering-Plough, and has been involved in performing clinical trials with many pharmaceutical industries that manufacture drugs used for the treatment of AE. SB has received research grants from La Fondation pour la Dermatite Atopique and Pierre Fabre Laboratories, has received personal fees from Bioderma, Ferring, La Roche Posay Laboratoire Dermatologique, Novalac and Sanofi-Genzyme, and has received non-financial support from Abbvie, Janssen and Novartis. SW has served as a consultant to Astellas, Novartis and Sanofi-Genzyme, has received independent research grants from Biogen, Novartis and Pfizer, and has been involved in performing clinical trials with many pharmaceutical industries that manufacture drugs used for the treatment of AE. CF has advised Roche/Genentech and Sanofi/Regeneron. The authors state there are no other conflicts of interests.

ACKNOWLEDGMENTS

We would like to acknowledge the Dutch atopic eczema patient society (VMCE, Vereniging voor Mensen met Constitutioneel Eczeem), the UK National Eczema Society and the Irish Skin Foundation for their support.

REFERENCES

- Beck LA, Thaci D, Hamilton JD, Graham NM, Bieber T, Rocklin R, et al. Dupilumab treatment in adults with moderate-to-severe atopic dermatitis. *N Engl J Med* 2014;371:130-9.
- Garcia-Doval I, Carretero G, Vanaclocha F, Ferrandiz C, Dauden E, Sanchez-Carazo JL, et al. Risk of serious adverse events associated with biologic and nonbiologic psoriasis systemic therapy: patients ineligible vs eligible for randomized controlled trials. *Arch Dermatol* 2012;148:463-70.
- Garritsen FM, Roekevisch E, van der Schaft J, Deinum J, Spuls PI, de Bruin-Weller MS. Ten years experience with oral immunosuppressive treatment in adult patients with atopic dermatitis in two academic centres. *J Eur Acad Dermatol Venereol* 2015;29:1905-12.
- Gerbens LA, Boyce AE, Wall D, Barbarot S, de Booi RJ, Deleuran M, et al. TREATment of ATopic eczema (TREAT) Registry Taskforce: protocol for an international Delphi exercise to identify a core set of domains and domain items for national atopic eczema registries. *Trials* 2017;18:87.
- Hay RJ, Johns NE, Williams HC, Bolliger IW, Dellavalle RP, Margolis DJ, et al. The global burden of skin disease in 2010: an analysis of the prevalence and impact of skin conditions. *J Invest Dermatol* 2014;134:1527-34.
- Proudfoot LE, Powell AM, Ayis S, Barbarot S, Baselga Torres E, Deleuran M, et al. The European TREATment of severe Atopic eczema in children Taskforce (TREAT) survey. *Br J Dermatol* 2013;169:901-9.
- Roekevisch E, Spuls PI, Kuester D, Limpens J, Schmitt J. Efficacy and safety of systemic treatments for moderate-to-severe atopic dermatitis: a systematic review. *J Allergy Clin Immunol* 2014;133:429-38.
- Simpson EL, Bieber T, Guttman-Yassky E, Beck LA, Blauvelt A, Cork MJ, et al. Two Phase 3 Trials of Dupilumab versus Placebo in Atopic Dermatitis. *N Engl J Med* 2016;375:2335-48.
- Spelsberg A, Prugger C, Doshi P, Ostrowski K, Witte T, Husgen D, et al. Contribution of industry funded post-marketing studies to drug safety: survey of notifications submitted to regulatory agencies. *BMJ* 2017;356:j337.
- Taylor K, Swan DJ, Affleck A, Flohr C, Reynolds NJ, DCTN. cwUTatU. Treatment of moderate-to-severe atopic eczema in adults within the UK: results of a national survey of dermatologists. *Br J Dermatol* 2016;In press.

Thaci D, Simpson EL, Beck LA, Bieber T, Blauvelt A, Papp K, et al. Efficacy and safety of dupilumab in adults with moderate-to-severe atopic dermatitis inadequately controlled by topical treatments: a randomised, placebo-controlled, dose-ranging phase 2b trial. *Lancet* 2016;387:40-52.

Totri CR, Eichenfield LF, Logan K, Proudfoot L, Schmitt J, Lara-Corrales I, et al. Prescribing practices for systemic agents in the treatment of severe pediatric atopic dermatitis in the US and Canada: The PeDRA TREAT survey. *J Am Acad Dermatol* 2017;76:281-5.

van der Schaft J, Politiek K, van den Reek JM, Christoffers WA, Kievit W, de Jong EM, et al. Drug survival for ciclosporin A in a long-term daily practice cohort of adult patients with atopic dermatitis. *Br J Dermatol* 2015;172:1621-7.

Williams H, Stewart A, von Mutius E, Cookson W, Anderson HR, International Study of A, et al. Is eczema really on the increase worldwide? *J Allergy Clin Immunol* 2008;121:947-54 e15.

Zaletel M, Kralj M. Methodological guidelines and recommendations for efficient and rational governance of patient registries. 2015.

TABLES

Table 1. Core set of domains and domain items to be captured in national atopic eczema registries*

DOMAINS	DOMAIN ITEMS
Demographics	Date of birth and date of enrolment into registry
	Gender
	Ethnicity
	Educational status
	Current occupation or education
AE diagnosis	How diagnosis AE is established
	Use of validated diagnostic criteria
	Date of onset AE
Past AE treatments	Previous phototherapy
	Previous systemic therapy
	Previous topical treatments for AE
	Previous day hospital care treatments for AE (outpatient)
	Previous hospitalization for AE
Current AE treatments	Phototherapy
	Systemic immunosuppressive therapy
	Topical treatments
	Amount of topical creams/ointments used per week
Family history of AE or allergic diseases	Family history of AE or allergic diseases
Allergic co-morbidities	Asthma
	Allergic rhinoconjunctivitis
	Atopic eye disease
	Eosinophilic oesophagitis
	Food allergies
	Contact allergies
Other co-morbidities	Past malignancies
	Past serious infections
Current concomitant medication (i.e. other than specific AE medication)	Antihistamines, oral or topical
	Topical antibiotics
	Oral antibiotics
	Immunosuppressives for other inflammatory diseases
General AE questions	Exposures that trigger disease flares
	Episodes of skin infection (i.e. folliculitis, HSV, molluscum contagiosum)
	Days lost from school/college/work
Baseline physical examination	Fitzpatrick skin type
	Skin examination
Baseline physician and patient reported domains	Physician-assessed clinical signs
	Investigator/physician global assessment
	Patient-reported symptoms
	Patient global assessment
	Generic quality of life score
	Skin-specific quality of life score
	Patient-reported satisfaction with AE care received
	Impact of AE on the family
Baseline investigations and assessments	Medical history (tuberculosis, HIV, hepatitis B or C)

	Full blood count
	Liver function
	Kidney profile
	Evaluating TPMT level prior to azathioprine use
Baseline management	Reasons for choosing specific treatment (systemic or phototherapy)
	Routine recording of relative contraindication(s) for selected treatment
Follow up general AE questions	Date of death and relation to AE
	Change in diagnosis after enrolment (e.g. from AE to CTCL)
Follow up serious adverse events	Serious adverse events
Follow up adverse events that cause stop or switch of therapy or change in dosage	Adverse events that cause stop or switch of therapy or change in dosage
<i>For follow up (serious) adverse events: probability of relationship with treatment</i>	<i>For (serious) adverse events: probability of relationship with treatment</i>
Follow up physical examination	Skin examination
Follow up physician and patient reported domains	Physician-assessed clinical signs
	Investigator/physician global assessment
	Patient-reported symptoms
	Patient global assessment
	Generic quality of life score
	Skin-specific quality of life score
	Reporting of disease control, e.g. flares, fully controlled weeks
	Adherence to treatment between appointments
	Patient-reported satisfaction with AE care received
	Impact of AE on the family
Follow up management	Reason for switching therapy
	Reason for discontinuation of therapy

Abbreviations: AE, atopic eczema; CTCL, cutaneous T cell lymphoma; HSV, herpes simplex virus; HIV, human immunodeficiency virus, TPMT, thiopurine methyltransferase.

* Results may slightly change after the upcoming 'How' to measure meeting.